

LAWOUJÒL

SE PA JIS YON TI GRATÈL



Lawoujòl ka danjere, sitou pou tibebe ak timoun piti.

Sentòm lawoujòl yo tipikman gen ladan yo:




Gwo lafyèv

(ka monte rive plis pase 104°F)



Tous



Nen kap koule



Wouj ak/oswa je ki plen dlo



Gratèl

(eklate

3-5 jou apre sentòm yo kòmanse)

Lawoujòl ka grav.

Lawoujòl ka lakòz konplikasyon sante grav, tankou nemoni, anflamasyon nan sèvo a (ansefalit) ak lanmò.



1 sou 5 moun ki pran lawoujòl pral lopitalize.



1 sou chak 20 timoun ki gen lawoujòl ap pran nemoni, ki se kòz ki pi komen pou lanmò akòz lawoujòl lakay timoun piti yo.



1 sou chak 1,000 moun ki gen lawoujòl ap devlope anflamasyon nan sèvo, sa ki ka lakòz domaj nan sèvo.



1 a 3 sou chak 1,000 moun ki gen lawoujòl ap mouri.

Lawoujòl kontajye anpil

epi li gaye nan lè a lè yon moun ki enfekte touse oswa estènye. Yon moun ki gen lawoujòl ka pase l bay 9 sou 10 moun ki pa pran vaksen bò kote yo.

Vaksinasyon se pi bon fason pou prevni lawoujòl

Yon timoun ta dwe pran vaksen kont lawoujòl-malmouton-ribeyòl (MMR) a laj 12-15 mwa ak 4-6 an. De dòz yo efikas 97% kont lawoujòl.

Klinik .vaksinasyon .CDPH .yo .bay .

Vaccines for Children
Iowa HHS



Vaccines for Adults
Iowa HHS



**IOWA
MEDICAL
SOCIETY**

Original content created by CDPH

